

- TOWNSEND, C. H. T. 1902. Contribution to a knowledge of the Coleopterous fauna of the lower Rio Grande valley in Texas and Tamaulipas, with biological notes and special reference to geographical distribution. *Trans. Texas Acad. Sci.* 5:49-101.
- TYSON, W. H. 1970. Notes on the host, larval habits, and parasites of two Texas cerambycids. *Proc. Ent. Soc. Wash.* 72:93.
- VOGT, G. B. 1949. Notes on Cerambycidae from the lower Rio Grande valley, Texas. *Pan-Pac. Ent.* 25:137-144; 175-184.
- WICKHAM, H. F. 1898. Recollections of old collecting grounds, pt. III, the lower Rio Grande valley (cont.). *Ent. News* 9:39-41.



TECHNIQUES

A new approach to measuring beetles was recently developed by me, based on observations of techniques used by mammalogists (craniometer). Using a Brinkmann two-axis stage as a specimen transport device, I had Spaulding Instruments of Pasadena, California, develop a potentiometer coupler to a Brinkmann-adapted micrometer head. The potentiometer is wired to a Spaulding Digital read-out box (x and y axis). The microscope (any will do) has an ocular cross-hair, and is mounted on a botanical scope swing-arm to allow room for the stage. The cross-hair aligns point A on the specimen, the digital read-out is "zeroed," the transport of the specimen to point B is viewed through the scope, and cross-hair alignment is set at B. Meanwhile, the potentiometer records electronically the movement of the micrometer head and relays the impulses to the digital device, recording the movement to 1/1000 of a millimeter. The reading can be taken by hand, keyed into a calculator, or transmitted to a computer via hook-up to an office terminal. The advantages of this system are repeatable results by multiple users (no ocular micrometer paralax problems), speed of measuring, resolution of measurements, and direct hook-up to a computer for immediate analysis of measurements.

-T. L. Erwin

